

Accepted by *The Astrophysical Journal Letters*

# Probing Population III Stars in Galaxy IOK-1 at $z = 6.96$ through He II Emission

Zheng Cai<sup>1</sup>, Xiaohui Fan, Linhua Jiang, Fuyan Bian, Ian McGreer, Romeel Davé, Eiichi Egami, Ann Zabludoff

*Steward Observatory, University of Arizona, Tucson, AZ 85721*

Yujin Yang

*Max-Planck-Institut für Astronomie, Königstuhl 17, D- 69117, Heidelberg, Germany*

S. Peng Oh

*Department of Physics, University of California, Santa Barbara, CA 93106*

## ABSTRACT

The He II  $\lambda 1640$  emission line has been suggested as a direct probe of Population III (Pop III) stars at high-redshift, since it can arise from highly energetic ionizing photons associated with hot, metal free stars. We use the HST WFC3/F130N IR narrowband filter to probe He II  $\lambda 1640$  emission in galaxy IOK-1 at  $z = 6.96$ . The sensitivity of this measurement is  $\gtrsim 5\times$  deeper than for previous measurements. From this deep narrowband imaging, combined with broadband observations in the F125W and F160W filters, we find the He II flux to be  $1.2 \pm 1.0 \times 10^{-18}$  ergs s<sup>-1</sup> cm<sup>-2</sup>, corresponding to a  $1\sigma$  upper limit on the Pop III star formation rate (SFR) of  $\sim 0.5$  M<sub>⊙</sub> yr<sup>-1</sup> for the case of a Salpeter IMF with  $50 \lesssim M/M_{\odot} \lesssim 1000$  and mass loss. Given that the broadband measurements can be fit with a UV continuum spectral flux density of  $\sim 4.85 \times 10^{-10} \times \lambda^{-2.46}$  ergs s<sup>-1</sup> cm<sup>-2</sup> Å<sup>-1</sup>, which corresponds to an overall SFR of  $\sim 16_{-2.6}^{+2.6}$  M<sub>⊙</sub> yr<sup>-1</sup>, massive Pop III stars represent  $\lesssim 6\%$  of the total star formation. This measurement places the strongest limit yet on metal-free star formation at high redshift, although the exact conversion from He II luminosity to Pop III SFR is highly uncertain due to the unknown IMF, stellar evolution, and photoionization effects. Although we have not detected He II  $\lambda 1640$  at more than the  $1.2\sigma$  level, our work suggests that a  $\gtrsim 3\sigma$  level detection is possible with JWST.

---

<sup>1</sup>Email: caiz at email.arizona.edu

## 1. Introduction

The first stars, i.e., Population III stars with zero metallicity, are likely to be very massive, due to inefficient cooling and the large Jeans mass in the early Universe (Abel et al. 2002; Yoshida et al. 2006). Massive Pop III stars are probably key contributors both of the ionization photons that caused cosmic reionization and to the early phases of cosmic chemical evolution (Whalen et al. 2008; Mackey et al. 2003; Bromm et al. 2003). Although recent observations have provided major breakthroughs in understanding the universe at  $z \gtrsim 7$  (e.g., Bouwens et al. 2010a; Yan et al. 2011), detection of high redshift galaxies that contain Pop III stars remains a major challenge observationally. Indirect constraints on Pop III stars come from mapping the reionization history (Fan, Carilli, & Keating 2006) or from studying the chemical evolution and the enrichment history through the observation of Extremely Metal-Poor stars and Hyper Metal-Poor stars (Beers & Christlieb 2005; Bessel et al. 2004). There are still no direct detections of high redshift galaxies hosting star formation in zero metallicity gas.

Models show that Pop III stars are considerably hotter than present-day stars, for a given stellar mass (e.g., Johnson 2010; Bromm et al. 2003). Emission of ionizing radiation, specifically photons with energies above 54.4eV that ionize He II, is enhanced by the high surface temperature photospheres of Pop III stars. He II  $\lambda 1640$  has been long suggested as a direct probe of Pop III stars (Tumlinson & Shull 2000; Oh 2001; Schaerer 2002). HeII  $\lambda 1640$  emission is also relatively easy to model compared to resonance lines such as Ly $\alpha$  or He II  $\lambda 304$ . The optical depth to He II  $\lambda 1640$  is small in almost all circumstances, so it does not suffer the radiative transfer effects that complicate the interpretation of Ly $\alpha$  flux in high-redshift galaxies (Schaerer 2003). In addition, it does not suffer from IGM Gunn-Peterson absorption as Ly $\alpha$  does (Tumlinson & Shull 2000).

How can He II  $\lambda 1640$  emission from star formation in zero metallicity gas be distinguished from other sources? Calculations show that the He II  $\lambda 1640$  emission in a metal-poor stellar population with metallicity as low as  $Z \sim 10^{-3}$  is still  $< 10^{-3}$  times lower than that in a population formed in zero metallicity gas (Tumlinson et al. 2003; Schaerer 2003). It is also known that nebular He II  $\lambda 4686$  was detected in a fairly large fraction of metal-poor HII regions (e.g., Skillman & Kennicutt 1993). The follow-up observations and theoretical investigations suggest that Wolf-Rayet (W-R) stars are responsible for this nebular He II emission (de Mello et al. 1998), but the hardness of this emission more than one order of magnitude weaker than that originated by massive Pop III stars (Schaerer 2003; Jimenez & Haiman 2006). Also, the He II  $\lambda 1640$  equivalent width of massive Pop III stars could be a factor of 2 larger than that of the typical AGN (Prescott et al. 2009; Elvis et al. 1994).

A number of studies have been carried out to search for He II emission in high-redshift

galaxies at  $4 < z < 6.6$ . Dawson et al. (2004), Ouchi et al. (2008), and Nagao et al. (2005) searched for He II  $\lambda 1640$  in either stacked or individual spectra of Ly $\alpha$  emitting galaxies. Nagao et al. (2008) carried out a survey for Ly $\alpha$ -He II emitters using a combination of intermediate and narrow-band filters in the optical window. But these observations so far have yielded only nondetections, constraining the massive Pop III star formation rate (hereafter  $\text{SFR}_{\text{PopIII}}$ ) to a few  $\text{M}_{\odot} \text{ yr}^{-1}$ , usually a few tenths of the overall SFR in these galaxies. Yet Jimenez & Haiman (2006) predict  $\text{SFR}_{\text{PopIII}}$  still to be significant at  $z \sim 3\text{--}5$ . Even stronger metal-free star formation is expected at higher redshifts, due to less chemical feedback from Pop III stars after their initial burst. However, at  $z > 7$ , He II is in the J-band, making ground-based observations extremely challenging. Due to its large field of view, high throughput, and low background, the new IR channel of the Wide Field Camera 3 (WFC3) on HST is the most powerful tool for detecting galaxies at  $z \gtrsim 7$  and for studying galaxy evolution at the end of cosmic reionization.

In this paper, we report the strongest upper limit yet on the contribution of Pop III stars in high-redshift galaxies, using deep narrow band imaging of galaxy IOK-1 (Iye et al. 2006) at  $z = 6.96$  in the HST/WFC3 F130N band, which includes He II  $\lambda 1640$  at this redshift. We have also carried out deep observations in broad bands to measure the continuum level and UV-based total SFR. In §2, we discuss the observations and data reduction. Photometric results, including the limit on He II flux, are presented in §3. We discuss the implication of our results on Pop III star formation in IOK-1 in §4. Throughout this paper, we adopt a cosmology based on the fifth year Wilkinson Microwave Anisotropy Probe (WMAP) data (Komatsu 2009):  $\Omega_{\Lambda} = 0.72$ ,  $\Omega_m = 0.28$ ,  $\Omega_b = 0.046$ , and  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ .

## 2. Observation and the data reduction

Our target, IOK-1, was discovered in a narrow band Ly $\alpha$  emitter survey using the NB973 filter on the Subaru Suprime-Cam (Iye et al. 2006), which is centered on Ly $\alpha$  at  $z \simeq 7.02$ . This galaxy is spectroscopically confirmed at  $z = 6.96$ . IOK-1 has a total Ly $\alpha$  luminosity of  $1.1 \pm 0.2 \times 10^{43} \text{ erg s}^{-1}$ , corresponding to a Ly $\alpha$ -based SFR  $\text{SFR}_{\text{Ly}\alpha} \sim 10 \pm 2 \text{ M}_{\odot} \text{ yr}^{-1}$ . (Iye et al. 2006).<sup>1</sup>

In the observed frame, the He II  $\lambda 1640$  emitted by IOK-1 is located at  $13054 \text{ \AA}$ , which is in the passband of the WFC3 F130N narrow band filter ( $\lambda_0 = 13006 \text{ \AA}$  and FWHM =

---

<sup>1</sup>These error bars only reflect the photometric uncertainties. There are big systematic uncertainties due to Ly $\alpha$  radiative transfer effect, galaxy IMF and metallicity assumptions. The  $\text{SFR}_{\text{Ly}\alpha}$  could change by a factor of 2 given these systematic uncertainties (also see §4).

150 Å). Resonant scattering of Ly $\alpha$  photons and IGM absorption of the blue wing usually systematically redshift the Ly $\alpha$  line center up to  $\sim 700 \text{ km s}^{-1}$  against the systemic velocity (e.g., Steidel et al. 2010). Given that He II will be located slightly blueward of the peak inferred from Ly $\alpha$  peak and have a narrower line width, we expect that the entire He II line falls securely into the most sensitive part of the F130N filter. IOK-1 was observed by HST WFC3 in March 2010. Eight orbits ( $\sim 20,000$  sec integration) were devoted to the F130N filter to measure the He II flux; we also observed this galaxy using the F125W filter for two orbits to determine the underlying continuum level and UV continuum-based SFR. A F160W band observation was also carried out (Egami et al. 2011, in preparation).

The F130N observations are divided into two visits, each consisting of a 4-point dither sequence with one-orbit per dither. During the the first visit, the first two dither positions were affected by the presence of a bright ghost image very close to IOK-1. We exclude these two images from further analysis. The F125W continuum observation was a two-orbit single visit, which is divided into two identical 4-point dither sequences, one per orbit. No ghost image affected the F125W observations. The individual images were reduced by WFC3-IR standard pipelines. Both the F130N and F125W images were combined using Multidrizzle (Koekemoer et al. 2002) with `final_scale` = 0.06, which is 0.48 of the original pixel size, and `final_pixfrac` = 0.7. High resolution images from the F130N and F125W bands are shown in Figure 1.

### 3. Results

#### 3.1. Photometry

Photometry is performed with SExtractor (Bertin & Arnouts 1996) using the rms map converted from the inverse variance image generated by Multidrizzle (Casertano et al. 2000). We measure the fluxes in the broadband (F125W and F160W) and the narrowband (F130N) images using the same Kron-like elliptical aperture determined from the F130N image (black elliptical aperture in Figure 1). The results are listed in Table 1. We also measure the flux in the F160W image of IOK-1 (Egami et al. 2011, in preparation), using the same aperture as used in the F125W and F130N images, finding a flux density of  $f_{\text{F160W}}^{\lambda} = 2.44 \pm 0.14 \times 10^{-20} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$ , corresponding to a magnitude of  $m_{\text{F160W}} = 25.98 \pm 0.06$ . We fit the photometry by assuming a model spectrum with a power law continuum with a dimensionless flux density  $f_{\text{con}} = A(\lambda/1\text{Å})^{\beta}$ , where A is a constant, and a narrow Gaussian for the He II emission line of flux  $F_{\text{HeII}}$ . The best-fit SED and photometry are shown in Figure 2. We find:

$$f_{\text{con}}(\lambda) = (4.85 \pm 0.23) \times 10^{-10} (\lambda/1 \text{ Å})^{-2.46 \pm 0.36}, \quad (1)$$

$$F_{\text{HeII}} = (1.2 \pm 1.0) \times 10^{-18} \text{ ergs s}^{-1} \text{ cm}^{-2}. \quad (2)$$

At  $z = 6.96$ , this corresponds to a total HeII luminosity of  $L_{\text{HeII}} = F_{\text{HeII}} \times 4\pi D_L^2 = (6.6 \pm 5.5) \times 10^{41} \text{ ergs s}^{-1}$ , or a  $2\text{-}\sigma$  upper limit of  $1.11 \times 10^{42} \text{ erg s}^{-1}$ . This  $2\text{-}\sigma$  upper limit is a factor of  $\gtrsim 4\text{--}5$  deeper than previous measurements. The rest-frame equivalent width of He II is  $4.2 \pm 3.5 \text{ \AA}$ .<sup>2</sup> Our results also show that IOK-1 has a blue continuum. Bouwens et al. (2010b) studied the value of the UV-continuum slope  $\beta$  in the Hubble Ultra Deep Field (HUDF). For luminous  $L_{z=3}^*$  galaxies (Steidel et al. 1999),  $\beta \sim -2.0 \pm 0.2$ . For lower luminosity  $0.1L_{z=3}^*$  galaxies,  $\beta \sim -3.0 \pm 0.2$ . IOK-1 has an absolute magnitude  $M_{\text{AB}}(1500\text{\AA}) = -21.3$ , comparable to  $L_{z=3}^*$ .

### 3.2. Morphology

The half-light radius  $r_{1/2}$  of IOK-1, based on our SExtractor measurements (Bertin & Arnouts 1996) and corrected for WFC3 PSF broadening, is  $0''.12$  in F125W. This corresponds to  $0.62 \pm 0.04 \text{ kpc}$ . IOK-1 is an extremely compact galaxy, consistent with the size of  $L_{z=3}^*$  galaxies at  $z \sim 7$  detected in the HUDF09 (Oesch et al. 2010). The observed surface brightness of IOK-1 is  $\mu_J \sim 24.5 \text{ mag arcsec}^{-2}$ , corresponding to a surface brightness of  $\mu_{rest} \sim 15.5 \text{ mag arcsec}^{-2}$ , after  $(1+z)^4$  correction for cosmological dimming. From Figure 1, it is obvious that IOK-1 consists of two components. We use GALFIT (Peng et al. 2002) to deblend the F125W continuum image, assuming an exponential profile for both components. The F125W image shows roughly equal brightness for the two components. The north west component has an effective radius of  $0.49 \pm 0.04 \text{ kpc}$ ; and the southeast component has an effective radius of  $0.57 \pm 0.03 \text{ kpc}$ . The two components are projected about  $0''.2$  ( $\sim 1 \text{ kpc}$ ) away from each other. More detailed discussions on the morphology are presented in

---

<sup>2</sup>Note that our observations could only detect He II flux stronger than  $3 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2}$  at  $3\sigma$  level, corresponding to a He II  $\lambda 1640$  equivalent width of  $\sim 11 \text{ \AA}$ , which is a factor of  $\gtrsim 6$  larger than that of normal stellar populations in the Lyman break spectrum of stacked  $z \sim 3$  galaxies (Shapley et al. 2003), and a factor of  $\gtrsim 3$  greater than the simulated maximum He II equivalent width generated by W-R stars (Brinchmann et al. 2008).

Table 1: SExtractor photometry in different bands

Filter	Flux Density $f^\lambda$ ( $10^{-20} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$ )	$m_{\text{AB}}$
F130N	$4.39 \pm 0.68$	$25.42 \pm 0.17$
F125W	$4.07 \pm 0.19$	$25.59 \pm 0.05$

Egami et al. (2011, in preparation). We also deblend the F130N narrow band image, which contains He II  $\lambda 1640$  emission. The north-west component is somewhat brighter. However, given the low signal-to-noise ratio of the data, this difference, while intriguing, is not statistically significant.

#### 4. Discussion

What limit can our observations place on the  $\text{SFR}_{\text{PopIII}}$  and its contribution to the overall star formation in IOK-1 at  $z \sim 7$ ? Photoionization by AGN or hot dense stellar winds from Wolf-Rayet stars (see e.g., Shapley et al. 2003; Brinchmann et al. 2008) may contribute to He II  $\lambda 1640$  emission although at a lower level than metal-free stars. More importantly, massive stars with the lowest, but non-zero metallicity ( $< 10^{-8}$ ) could generate a comparable amount of He II  $\lambda 1640$  emission to Pop III stars (Schaerer 2003). However, here we assume that the observed He II emission originates entirely from metal-free stars. Therefore, our derived  $\text{SFR}_{\text{PopIII}}$  should be regarded as an upper limit. In the case of constant star formation, at equilibrium, recombination line luminosities  $L_l$  are proportional to the  $\text{SFR}_{\text{PopIII}}$  (Schaerer 2002), so

$$\begin{aligned} L_{\text{HeII}} &= c_{1640}(1 - f_{\text{esc}})Q(\text{He}^+) \left( \frac{\text{SFR}_{\text{PopIII}}}{\text{M}_{\odot}\text{yr}^{-1}} \right) \\ &= L_{1640,\text{norm}} \left( \frac{\text{SFR}_{\text{PopIII}}}{\text{M}_{\odot}\text{yr}^{-1}} \right) \end{aligned} \quad (3)$$

where  $L_{1640,\text{norm}} \equiv c_{1640}(1 - f_{\text{esc}})Q(\text{He}^+)$  is the theoretical He II  $\lambda 1640$  line luminosity normalised to  $\text{SFR}=1 \text{ M}_{\odot} \text{ yr}^{-1}$ .  $c_{1640}$  is the He II  $\lambda 1640$  emission coefficient given in Table 1 of Schaerer (2003):  $c_{1640} = 5.67 \times 10^{-12} \text{ ergs}$  for  $T_e = 30\text{kK}$ , where nebular emission is calculated assuming case B recombination.  $Q(\text{He}^+)$  is the number of  $\text{He}^+$  ionizing photons per second,  $f_{\text{esc}}$  represents the fraction of total ionizing radiation released into the IGM without being coupled to the ISM in the galaxy. We assume  $f_{\text{esc}} = 0$ , as it is expected to be small among high-redshift galaxies (e.g., Gnedin et al. 2008). Schaerer (2002) calculates the total  $\text{He}^+$  ionizing photon flux for a Salpeter IMF with a range of lower and upper mass cut-offs and different assumptions of mass loss.

The conversion from He II  $\lambda 1640$  luminosity ( $L_{\text{HeII}}$ ) to  $\text{SFR}_{\text{PopIII}}$  is affected by a number of uncertainties, including: (1) IMF, (2) mass loss, and (3) photo-ionization model. For the IMF, theoretical calculations show that the lack of efficient cooling mechanisms generally result in an extremely top-heavy IMF compared to a galactic IMF (Abel et al. 2002; Bromm & Larson 2004; Yoshida et al. 2006; O’Shea & Norman 2006). However, both the shape and mass range of such an IMF are poorly constrained. Scannapieco et al. (2006)

Table 2: Pop III star formation rate

IMF(Salpeter)	Mass Loss	$L_{1640,\text{norm}}$ (ergs s <sup>-1</sup> )	SFR <sub>PopIII</sub> (M <sub>⊙</sub> yr <sup>-1</sup> )
$1 \lesssim M/M_{\odot} \lesssim 500$	No	$9.66 \times 10^{40}$	$6.8 \pm 4.7$
$50 \lesssim M/M_{\odot} \lesssim 500$	No	$6.01 \times 10^{41}$	$1.1 \pm 0.9$
$1 \lesssim M/M_{\odot} \lesssim 500$	Yes	$3.12 \times 10^{41}$	$2.1 \pm 1.5$
$50 \lesssim M/M_{\odot} \lesssim 1000$	Yes	$2.33 \times 10^{42}$	$0.3 \pm 0.2$

suggests a lower mass cutoff of  $0.8 M_{\odot}$  based on metal abundances in Galactic halo stars. Numerical simulations suggest that stars as massive as  $500 M_{\odot}$  can be formed by accretion into a primordial protostar (Bromm & Larson 2004), which is generally taken as an upper limit in different simulations (Schaerer 2002; Scannapieco et al. 2003; Tumlinson 2006; Raiter et al. 2010). Both a Salpeter IMF (Schaerer 2003; Scannapieco et al. 2003) and a Log-normal IMF (Tumlinson 2006) have been used in model calculations. Table 2 shows a factor of six difference in inferred SFR arising from different lower mass cutoffs in the Salpeter IMF.

As shown in Table 2, models with strong mass loss have a larger conversion factor  $L_{1640,\text{norm}}$ ; therefore, for the same HeII luminosity, they yield a lower SFR<sub>PopIII</sub> (Schaerer 2003). Kudritzki (2002) shows that very low-metallicity stars close to the Eddington limit are subject to non-zero mass loss. Smith & Owocki (2006) argue that massive shells around luminous blue variables and so-called supernova impostors indicate that continuum-driven winds or hydrodynamic explosions dominate the mass loss of very massive stars, a claim which is insensitive to metallicity and so could apply to Pop III stars. Ekstrom (2007) explores the effects of rotation, anisotropic mass loss, and magnetic fields on the core size of a Pop III star, pointing out that, under certain conditions, very massive stars ( $140M_{\odot} < M < 260M_{\odot}$ ) losing mass through rotation could avoid ending their lives as pair-instability supernovae (PISN). Whether Pop III stars end their lives as PISN or not will affect the metal ejection efficiency and hence affect the SFR<sub>PopIII</sub>. From Table 2, stronger mass loss will lower SFR<sub>PopIII</sub> by a factor of more than three.

The derived SFR also depends on the photoionization model. Schaerer (2002) calculates nebular emission lines using standard case B recombination, in which the optical depth of the HII region is large. Eq. (3) is derived under this assumption. However, for low metallicity nebulae ionized by very hot Pop III stars, the case B predictions for line and continuum emission may have non-negligible deviations from real nebular astrophysics (Raiter et al. 2010). Therefore, the SFR<sub>PopIII</sub> upper limits derived from Eq. (3) then need to be qualitatively revised here. According to Raiter et al. (2010), for a given He II luminosity, SFR<sub>PopIII</sub> could be a few times higher, depending on the ionizing parameters, e.g.,



the hydrogen number density  $n(H)$ , inner radius of the nebula  $r_{in}$ , and hydrogen ionizing photon flux  $Q(H)$ .

Our upper limit on the He II emission combined with our detection of IOK-1 in the Spitzer IRAC 1 and 2 bands (Egami et al., in prep.) suggest that IOK-1 is dominated by stars of metallicity above the critical value ( $Z_{crit} \sim 5 \times 10^{-4} Z_{\odot}$ ), leading to a normal IMF (e.g., Bromm 2004). Using the relation given in Madau et al. (1998), the rest frame UV luminosity of the galaxy IOK-1 corresponds to an overall SFR  $\sim 20_{-0.9}^{+0.9} M_{\odot} \text{ yr}^{-1}$ . These error bars only reflect the uncertainties in the J-band magnitude. The true errors are larger as the conversion from UV luminosity to overall SFR is highly and systematically uncertain. In this conversion, Madau (1998) assumes solar metallicity; lower metallicities down to 0.0004 could lower the SFR by a factor of 2 (Schaerer 2003). Furthermore, the conversion assumes a Salpeter IMF with mass ranging from 0.1 to  $125 M_{\odot}$ , whereas assuming a Scalo IMF would double the derived overall SFR. The UV-based overall SFR which should be less obscured is somewhat higher than that based on the Ly $\alpha$  emission line ( $\sim 10 \pm 2 M_{\odot} \text{ yr}^{-1}$ ; Iye et al. 2006), but consistent given the uncertainties in both the stellar population models and the Ly $\alpha$  radiative transfer effects.

The conversion of UV luminosity to an overall SFR above assumes a conversion factor between UV luminosity and SFR for Pop I and Pop II stellar populations, without correcting for Pop III stars with a top-heavy IMF. Using the fiducial Pop III Salpeter IMF of 50 - 500  $M_{\odot}$  and no mass loss, we find that our derived upper limit of  $2.0 M_{\odot} \text{ yr}^{-1}$  for  $\text{SFR}_{\text{PopIII}}$  corresponds to an upper limit of  $1.2 \times 10^{-20} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$  in F125W flux density. Note that the observed flux density is  $4.07 \pm 0.19 \times 10^{-20} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$ . After correcting for the UV-luminosity contributed by Pop III stars, the overall SFR is  $\sim 16_{-2.6}^{+2.6} M_{\odot} \text{ yr}^{-1}$ . (As stated previously, this result could change a factor of 2 given the systematic uncertainties associated with the IMF and metallicity assumptions). Thus, IOK-1 is not dominated by very massive Pop III stars. It should be noted that low mass metal-free stars do not produce an appreciable amount of He II  $\lambda 1640$  emission. Therefore, our observations only constrain the amount of high-mass Pop III star formation in the galaxy and cannot be used to rule out the existence of low mass, low surface temperature Pop III stars (although, as discussed in §4, most theoretical calculations favor a top heavy IMF for Pop III). Trenti & Stiavelli (2009) show that the fraction of Pop III stars  $f_{\text{III}}(z)$  per dark matter halo at  $z \sim 10$  is only a few thousandths. Davé et al. (2006) and Tumlinson (2006) also suggest that  $f_{\text{III}}(z)$  in halos at  $z \sim 7$  is small:  $\lesssim 1\%$ . Taking into account the systemic uncertainties in the UV-based overall SFR, our observations indicate that the galaxy IOK-1 cannot be dominated by Pop III stars with very top heavy IMFs. For example, for a Salpeter IMF with  $50 \lesssim M/M_{\odot} \lesssim 500$  and no mass loss, the ratio of  $\text{SFR}_{\text{PopIII}}$  to the overall SFR is  $\lesssim 25\%$ . For a Salpeter IMF with  $50 \lesssim M/M_{\odot} \lesssim 1000$  and mass loss, the ratio is  $\lesssim 6\%$ .



Our deep HST narrow-band imaging places the strongest constraint yet on the  $\text{SFR}_{\text{PopIII}}$  in high-redshift galaxies. Although we have not detected the He II emission line in IOK-1 at more than  $1.2\sigma$ , this limit suggests that detection of He II emission from Pop III stars may be already within the reach of current facilities. Future facilities, especially JWST and GSMT, will be likely to probe down to the  $\gtrsim 3\sigma$  level and directly detect the signatures of the earliest star formation in the universe.

We thank the anonymous referee for insightful comments which have significantly improved the Letter. We thank Stefano Casertano, Daniel Schaerer, and Haojing Yan for useful discussions. Support for this work was provided by NASA through grant HST-GO-11587 from the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS5-26555. ZC, XF, LJ, FB and IM acknowledge support from NSF grant AST 08-06861, and a David and Lucile Packard Fellowship. AIZ acknowledges support from the NSF through grant AST-0908280 and from NASA through grant NNX10AD47G.

## REFERENCES

- Abel, T., Bryan, G. L., & Norman, M. L. 2002, *Science*, 295, 93
- Beers, T. C., & Christlieb, N. 2005, *ARA&A*, 43, 531
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
- Bessel, M. S., Christlieb, N., & Gustafsson, B. 2004, *ApJ*, 612, L61
- Bouwens, R. J., et al. 2010a, *ApJ*, 709, L133
- . 2010b, *ApJ*, 708, L69
- Brinchmann, J., Pettini, M., & Charlot, S. 2008, *MNRAS*, 385, 769
- Bromm, V., & Larson, R. B. 2004, *A&A*, 42, 79
- Bromm, V., Yoshida, N., & Hernquist, L. 2003, *ApJ*, 596, L135
- Casertano, S., et al. 2000, *AJ*, 120, 2747
- Davé, R., Finlator, K., & Oppenheimer, B. D. 2006, *MNRAS*, 370, 273
- Dawson, S., Rhoads, J., & Malhotra, S. 2004, *ApJ*, 617, 707

- de Mello, D. F., Schaerer, D., Heldmann, J., & Leitherer, C. 1998, *ApJ*, 507, 199
- Elvis, M., et al. 1994, *ApJS*, 95, 1
- Fan, X., Carilli, C. L., & Keating, B. 2006, *ARA&A*, 44, 415
- Gnedin, N. Y., Kravtsov, A. V., & Chen, H. 2008, *ApJ*, 672, 765
- Hirschi, R., Maeder, A., Meynet, G., Chiappini, C., & Ekström, S. 2007, in *EAS Publications Series*, Vol. 24, *EAS Publications Series*, ed. E. Emsellem, H. Wozniak, G. Massacrier, J.-F. Gonzalez, J. Devriendt, & N. Champavert, 263–268
- Iye, M., Ota, K., Kashikawa, N., & Furusawa, H. 2006, *Nature*, 443, 186188
- Jimenez, R., & Haiman, Z. 2006, *Nature*, 440, 501
- Johnson, J. L. 2010, *MNRAS*, 404, 1425
- Koekemoer, A. M., Fruchter, A. S., Hook, R. N., & Hack, W. 2002, in *The 2002 HST Calibration Workshop : Hubble after the Installation of the ACS and the NICMOS Cooling System*, ed. S. Arribas, A. Koekemoer, & B. Whitmore, 337–+
- Komatsu, E. e. a. 2009, *Astrophys.J.Suppl*, 540, 330
- Kudritzki, R. P. 2002, *ApJ*, 577, 389
- Mackey, J., Bromm, V., & Hernquist, L. 2003, *ApJ*, 586, 1
- Madau, P., Pozzetti, L., & Dickinson, M. 1998, *ApJ*, 498, 106
- Nagao, T., Maiolino, R., & Marconi, A. 2005, *ApJ*, 631, L5
- Nagao, T., Sasaki, S. S., & Maiolino, R. 2008, *ApJ*, 680, 100
- Oesch, P. A., et al. 2010, *ApJ*, 709, L21
- Oh, S. P. 2001, *ApJ*, 553, 499
- O’Shea, B. W., & Norman, M. L. 2006, *ApJ*, 648, 31
- Ouchi, M., Shimasaku, K., & Akiyama, M. 2008, *ApJS*, 176, 301
- Peng, C. Y., Ho, L. C., Impey, C. D., & Rix, H. 2002, *AJ*, 124, 266
- Prescott, M. K. M., Dey, A., & Jannuzi, B. T. 2009, *ApJ*, 702, 554

- Raiter, A., Schaerer, D., & Fosbury, R. 2010, ArXiv e-prints
- Scannapieco, E., Kawata, D., Brook, C. B., Schneider, R., Ferrara, A., & Gibson, B. K. 2006, *ApJ*, 653, 285
- Scannapieco, E., Schneider, R., & Ferrara, A. 2003, *ApJ*, 589, 35
- Schaerer, D. 2002, *A&A*, 382, 28
- . 2003, *A&A*, 397, 527
- Shapley, A. E., Steidel, C. C., Pettini, M., & Adelberger, K. L. 2003, *ApJ*, 588, 65
- Skillman, E. D., & Kennicutt, Jr., R. C. 1993, *ApJ*, 411, 655
- Smith, N., & Owocki, S. P. 2006, *ApJ*, 645, L45
- Steidel, C. C., Adelberger, K. L., Giavalisco, M., Dickinson, M., & Pettini, M. 1999, *ApJ*, 519, 1
- Steidel, C. C., Erb, D. K., Shapley, A. E., Pettini, M., Reddy, N., Bogosavljević, M., Rudie, G. C., & Rakic, O. 2010, *ApJ*, 717, 289
- Trenti, M., & Stiavelli, M. 2009, *ApJ*, 694, 879
- Tumlinson, J. 2006, *ApJ*, 641, 1
- Tumlinson, J., Michael, S. J., & Venkatesan, A. 2003, *ApJ*, 584, 608
- Tumlinson, J., & Shull, J. M. 2000, *ApJ*, 528, L65
- Whalen, D., O’Shea, B. W., Smidt, J., & Norman, M. L. 2008, *ApJ*, 679, 925
- Yan, H., Yan, L., Zamojski, M., Windhorst, R., McCarthy, P., Fan, X., & Rottgering, *ApJ*, 728, L22
- Yoshida, N., Omukai, K., Hernquist, L., & Abel, T. 2006, *ApJ*, 652, 6




Fig. 1.— High resolution images of galaxy IOK-1 in the F125W (left) and the F130N (right) bands, with contours spaced by 1.4 sky rms for the F125W image and 1 sky rms for the F130N image. The black elliptical aperture is determined by the F130N image. IOK-1 is clearly resolved into two components separated by  $\sim 0''.2$ . Photometric analysis of the F130N image reveals a flux density excess of  $1.2\sigma$ .




Fig. 2.— Best-fit spectrum (black line) of galaxy IOK-1 with the total UV continuum, as well as the  $\text{Ly}\alpha$  and He II emission lines. The filter response curves of F125W (blue dashed line), F160W (dark blue dashed line) and F130N (brown dashed line) are plotted. In addition, photometry in three different bands is overplotted at the effective wavelength of each filter.